

## Text of Agrispon Household Use Sublabel.

# AGRISPON®

A BIOLOGICALLY DERIVED  
BIOSTIMULANT FOR SOIL AND PLANTS

### ACTIVE INGREDIENT:

Plant Extract\* 0.56%

OTHER INGREDIENTS 99.44%

TOTAL 100.00%

\* The plant extract is derived from *Quercus falcata*, *Opuntia lindheimeri*, *Rhus aromatica*, and *Rhizophora mangle* tissues.

Keep out of reach of children

### CAUTION

See Attachment for FIRST AID, PRECAUTIONARY STATEMENTS, and DIRECTIONS FOR USE

Manufactured by:

Agriculture Sciences, Inc.

3227 Garden Brook EPA

Dallas, TX 75234

(972) 243-8930

www.agsciinc.com

EPA Reg. No. 59174-3

Est. No. 59174-TX-1

Agrispon stimulates root development and beneficial soil micro-organisms, increasing the efficiency of nutrient uptake from soil and improving soil structure over time. Agrispon can also hasten certain plant growth processes and improve a plant's ability to withstand pest and environmental stresses. Agrispon enables the soil to supply nutrients to the plant more efficiently.

### USES

Use Agrispon on all food and feed crops, consistent with the use directions and restrictions stated in this attachment. Also, use Agrispon on flowers and shrubs, as noted in this attachment.

### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

### FIRST AID

If on skin or clothing:	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
-------------------------	---

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call 1-800-274-8930, Monday through Friday, 9 AM to 5 PM (central time). After 5 PM (central time) call your poison control center at 1-800-222-1222.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Remove clothing/PPE immediately after handling this product. Wash the outside of gloves before removing.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate.

### AGRISPON DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use only on plants grown for other than commercial or research purposes, which may include plants in habitations, home fruit, and vegetable gardens, and home greenhouses.

Keep children and pets out of the treated area until sprays have dried.

### EQUIPMENT AND MIXING

Apply Agrispon via hose-end, hand - carried, or backpack spray equipment. Shake well before using. Agitate solutions either during or immediately after dilution. Apply solution within eight hours of mixing. Dilute Agrispon in water as specified in the table below.

Application Site	Treatment Area	Water (min.)*
<b>Permanent plants</b> Food crops, Fruit trees, Ornamental plants	50 sq. ft.	1 pint
	500 sq. ft.	1 quart
<b>90 to 180 day growth cycle plants</b> Food Crops, Ornamental plants	1,000 sq. ft.	2 quarts
	5,000 sq. ft.	1.5 gal
	1 acre	5 gal

\*The table specifies the minimum amount of water to be used; dilute Agrispon in a larger volume of water if desired. The amount of water used will vary according to equipment, type of nozzle used, number of nozzles, ground speed, system pressure and calibration. If soil is covered with plant material, mulch, or thatch, use sufficient water to transport Agrispon to the soil during application or lightly irrigate after application.

### SPECIFIC CROP APPLICATION INSTRUCTIONS

Crop	Rate of Agrispon and Time
<b>FRUIT &amp; GRAPES</b>	
Fruit & Grapes	1.5 fl. oz./1,000 sq. ft. every 4 months for developing plants
	1.5 fl. oz./1,000 sq. ft. at initiation of flowering for plants in production
	1.5 fl. oz./1,000 sq. ft. at the emergence of the fruit
<b>VEGETABLES</b>	
Peppers, Peas,	1 tbsp./gal. of water 8 days after germination or transplant
Beans, Melon,	1 tbsp./gal. of water at initiation of flowering
Tomato, Egg	1 tbsp./gal. of water at fruit initiation
Plant	1 tbsp./gal. of water monthly during production
Corn (Sweet)	3 tbsp./gal. of water at third leaf
	3 tbsp./gal. of water 18 days after initial application
Onion	2 tsp./gal. of water at 15 days after transplanting
	2 tsp./gal. of water at the beginning of bulb development (60 days after transplant)
Potato	1 tsp./gal. of water during planting onto seed pieces before covering in furrow
	1 tsp./gal. of water at 4-5 leaf stage
Carrot, Radishes	1 tsp./gal. of water at hilling
	1 tsp./gal. of water 8 days after germination or transplant
Brassica	1 tsp./gal. of water monthly during production
	1 fl. oz./gal. of water to foliage and soil at germination or seeding
Lettuce	1 fl. oz./gal. of water to foliage and soil at flowering
	1 fl. oz./gal. of water to foliage and soil at flower initiation
<b>FLOWERS AND SHRUBS</b>	1 fl. oz./gal. to foliage and soil 2-6 days after emergence
	1 fl. oz./gal. of water to foliage and soil one week after transplanting
Annual Flowers	2 tbsp./gal. of water to foliage and soil every 4 weeks during active growth
	2 tbsp./gal. of water to soil and foliage after planting
Perennial Flowers	2 tbsp./gal. to soil and foliage monthly during maturation
	1 tbsp./gal. to foliage at weeks 2, 4, 6, & 8 during production
Fern	1 fl. oz./gal. to foliage and soil 30 days after transplanting
	1 fl. oz./gal. to foliage and soil on emergence after dormancy
Rose	1 fl. oz./gal. to foliage and soil 90 days after initial treatment
	2 tbsp./gal. of water to foliage and soil every 15 days during continuous production
Non Flower Annuals	1 tbsp./gal. of water to foliage under conditions of stress*
	1 tbsp./gal. of water to foliage at week 1, 4 & 8 after pruning
Woody Ornamentals	2 tbsp./gal. of water to foliage and soil at weeks 2 & 4 during rooting
	2 tbsp./gal. of water to foliage 1 week before transplanting for liners
Woody Ornamentals	2 tbsp./gal. of water to foliage and soil 1 week after potting
	2 tbsp./gal. of water to foliage and soil every 15 days for cuttings
Woody Ornamentals	2 tbsp./gal. of water to foliage and soil at weeks 2 & 4 during rooting
	2 tbsp./gal. of water to foliage 1 week before transplanting for liners
Woody Ornamentals	2 tbsp./gal. of water to foliage and soil 1 week after potting
	2 tbsp./gal. of water to foliage and soil 1 week after potting

\*Stress from temperature, humidity, low light and phytotoxicity from agrichemicals.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

#### PESTICIDE STORAGE

Store in original container only.

#### CONTAINER DISPOSAL AND CONTAINER HANDLING

##### If empty:

Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

##### If partly filled:

Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

### WARRANTY

Manufacturer warrants that this product conforms to the original formulation and is fit for use as directed. To the extent consistent with applicable law, neither Manufacturer nor seller shall be liable for any injury, loss or damage, direct or indirect, arising from misuse of the product. To the extent consistent with applicable law, Agriculture Sciences, Inc. and its various sellers' only obligation shall be to replace such quantity of the product that is proven defective before purchase.

© 2012 Agriculture Sciences, Inc.