



# AGRISPON<sup>®</sup>

**BIOLOGICALLY DERIVED  
BIOSTIMULANT FOR SOIL AND PLANTS**

## **TURF & ORNAMENTALS**



### **FOR THE SOIL**

- Increases the efficiency of root development and absorption of nutrients
- Promotes development of microbial fauna
- Increases availability of nitrogen in crop residues and fertilizers
- Increases the development of roots and mycorrhizae
- Augments the activity of the rhizosphere
- Quicker breakdown of crop residues
- Helps loosen tight soils



### **FOR THE PLANT**

- Better development of stems and leaves
- Increases the production of fruit, grains, roots, tubers, pods, ears, racemes
- Augments chlorophyll production and rate of photosynthesis
- Stimulates growth, maturation and quality of fruit
- Better sugar and protein content in plants
- Increased tolerance to drought and temperature extremes
- Better tolerance to damage from herbicides and other agrichemicals



# AGRISPON APPLICATION

Use Agrispon on all food and feed crops as well as landscape shrubs, ornamental and flowering plants including ornamental grasses. There is no waiting period between application and harvesting.

Agrispon can be applied with virtually all amateur/home gardener equipment and professional grower equipment, including hose end sprayer, backpack, boom, fan jet, blower, trombone, and chemigation.

PLANT	RATE		TIME & METHOD
	Oz./Acre	Liter/Ha	
Golf Courses & Other Recreational Turf	13	1	<ul style="list-style-type: none"> <li>• Every 45 days during active growth</li> </ul>
Fern	20	1.5	<ul style="list-style-type: none"> <li>• To foliage and soil 30 days after transplanting</li> <li>• At planting to foliage and soil on emergence after dormancy</li> <li>• At planting to foliage and soil 90 days after initial treatment</li> </ul>

PLANT	DILUTION RATE IN WATER		APPLY MIXTURE TO FOLIAGE . . .
	Tbsp/Gal	ml/Liter	
Carnation	2	8	<ul style="list-style-type: none"> <li>• Monthly for cuttings</li> <li>• And soil one week after transplanting</li> <li>• Between pinching and formation of buds</li> </ul>
	1	4	
	1	4	
Chrysanthemum	2	8	<ul style="list-style-type: none"> <li>• And soil 1 week after pruning</li> <li>• And soil after planting</li> <li>• And foliage monthly during maturation</li> </ul>
	2	8	
	1	4	
Baby's Breath	2	8	<ul style="list-style-type: none"> <li>• And soil every 15 days for cuttings</li> <li>• And soil at weeks 2 &amp; 4 after root initiation</li> <li>• 1 week before transplanting for liners</li> <li>• And soil 1 week after transplanting, 1 week before setting of light and pre-flowering</li> </ul>
Rose	2	8	<ul style="list-style-type: none"> <li>• And soil every 15 days during continuous production</li> <li>• Under conditions of stress*</li> <li>• At weeks 1, 4 &amp; 8 after pruning</li> </ul>
	1	4	
	1	4	
Other Flowers	6.5	0.5	<ul style="list-style-type: none"> <li>• And soil 1 week after transplant and in pre-flowering stage</li> </ul>
Annuals	13	1	<ul style="list-style-type: none"> <li>• And soil 2 &amp; 4 weeks after sticking</li> <li>• 1 week before transplanting for liners</li> <li>• And soil 1 week after potting</li> </ul>
Woody Ornamentals	2	8	<ul style="list-style-type: none"> <li>• And soil every 15 days for cuttings</li> <li>• And soil at 2 &amp; 4 weeks after root initiation</li> <li>• 1 week before transplanting for liners</li> <li>• And soil 1 week after potting</li> </ul>

\*Stress from temperature, humidity, low light and phytotoxicity from agrichemicals



www.agsciinc.com  
 info@agsciinc.com  
 800-274-8930

