



# AGRISPON<sup>®</sup>

BIOLOGICALLY DERIVED  
BIOSTIMULANT FOR SOIL AND PLANTS

**FRUIT, VEGETABLE & ORCHARD**



## FOR THE SOIL

- Increases the efficiency of root development and absorption of nutrients
- Promotes development of microbial fauna
- Increases availability of nitrogen in crop residues and fertilizers
- Increases the development of roots and mycorrhizae
- Augments the activity of the rhizosphere
- Quicker breakdown of crop residues
- Helps loosen tight soils



## FOR THE PLANT

- Better development of stems and leaves
- Increases the production of fruit, grains, roots, tubers, pods, ears, racemes
- Augments chlorophyll production and rate of photosynthesis
- Stimulates growth, maturation and quality of fruit
- Better sugar and protein content in plants
- Increased tolerance to drought and temperature extremes
- Better tolerance to damage from herbicides and other agrichemicals



# AGRISPON APPLICATION

Use Agrispon on all food and feed crops as well as landscape shrubs, ornamental and flowering plants including ornamental grasses. There is no waiting period between application and harvesting.

Agrispon can be applied with virtually all amateur/home gardener equipment and professional grower equipment, including hose end sprayer, backpack, boom, fan jet, blower, trombone, and chemigation.

CROP	RATE		TIME & METHOD
	Oz./Acre	Liter/Ha	
Brassica	18	1.3	<ul style="list-style-type: none"> <li>To foliage and soil at germination or seeding</li> <li>To foliage and soil at flowering</li> <li>To foliage and soil at flower initiation</li> </ul>
Carrot, Radishes	6.5	0.5	<ul style="list-style-type: none"> <li>8 Days after germination or transplant</li> <li>To foliage and soil at flowering</li> <li>Monthly during production</li> </ul>
Citrus	18	1.3	<ul style="list-style-type: none"> <li>Pre-flowering</li> <li>Halfway through production</li> <li>After harvest</li> </ul>
Coffee	11	0.8	<ul style="list-style-type: none"> <li>At flower initiation</li> <li>At start of fruit development</li> <li>After harvest</li> </ul>
Grapes & Other Fruit	6.5	0.5	<ul style="list-style-type: none"> <li>Every 4 months for developing plants</li> <li>At initiation of flowering for plants in production</li> <li>At the emergence of the fruit</li> </ul>
Lettuce	18	1.3	<ul style="list-style-type: none"> <li>To foliage and soil 2 - 6 days after emergence</li> </ul>
Onion	6.5	0.5	<ul style="list-style-type: none"> <li>At 15 days after transplanting</li> <li>At the beginning of bulb development (60 days after transplant)</li> </ul>
Peppers, Peas, Beans, Melon, Tomato, Egg Plant	6.5	0.5	<ul style="list-style-type: none"> <li>8 Days after germination or transplant</li> <li>At initiation of flowering</li> <li>At fruit initiation</li> <li>Monthly during production</li> </ul>
Potato	6.5	0.5	<ul style="list-style-type: none"> <li>During planting onto seed pieces before covering in furrow</li> <li>At 4 -5 leaf stage</li> </ul>
	13	1	<ul style="list-style-type: none"> <li>At hilling</li> </ul>



**AgSci**<sup>®</sup>

www.agsciinc.com

info@agsciinc.com

800-274-8930

